

HOVERFLY FAUNA (DIPTERA: SYRPHIDAE) OF THE LANDSCAPE OF OUTSTANDING FEATURES “VLASINA”

TAMARA TOT¹, MIHAILO VUJIĆ², LAURA LIKOV¹, ZORICA NEDELJKOVIĆ³,
SNEŽANA RADENKOVIĆ¹ and ANTE VUJIĆ¹

¹ University of Novi Sad, Faculty of Sciences, Department of Biology and Ecology,
Trg Dositeja Obradovića 2, Novi Sad 21000, Serbia
Email: tamaratot90@gmail.com, laura.likov@dbe.uns.ac.rs, snezana.radenkovic@dbe.uns.ac.rs,
ante.vujic@dbe.uns.ac.rs

² Bulevar revolucije 33, Vrčin, Beograd 11000, Serbia
Email: mihailovujic01@gmail.com

³ Biosense Institute-Research Institute for Information Technologies in Biosystems,
Dr Zorana Đinđića 1, Novi Sad 21000, Serbia
Email: zoricaned14@gmail.com

Abstract

In the many years of research organized between 1977 and 2017, a total of 76 hoverfly species were collected in the protected Landscape of Outstanding Features “Vlasina” (southeastern Serbia) on six localities. Among the identified taxa there is a rare species, *Cheilosia longula* (Zetterstedt, 1838), which is protected in Serbia. Additionally, one potentially new hoverfly species is registered (*Merodon* aff. *cinereus*). The first species checklist of hoverflies for Vlasina is proposed.

KEY WORDS: Syrphidae, Vlasina, fauna, new records, threatened species

Introduction

Hoverflies (Diptera: Syrphidae) represent one of the most species-rich dipteran families, with over 6000 described species from 188 genera (Thompson, 2013). European syrphid fauna accounts for about 870 species (Speight, 2016). Widespread in the world, they represent an ecologically important group of insects that perform ecosystem services such as plant pollination, predation of plant pests and nutrient recycling (Rotheray & Gilbert, 2011).

The Landscape of Outstanding Features “Vlasina” is located in southeastern Serbia, mostly in the municipality of Surdulica. The specificity of the Vlasina biotope is determined by the high diversity of its flora, fauna and ecosystems. Despite its relatively small area (127 km²), Vlasina has many endemic and protected species due to the presence of different types of habitats (mires, wetlands, forests, meadows and grasslands) (according to the Law on the Environmental Protection, Official Gazette of Republic of Serbia, No.66/91). The available data on the specific fauna structure of Vlasina refer mostly to vertebrate species, while data on insect fauna are scarce. In total, 11 species of Odonata, 42 species of Orthoptera (Pavićević & Zatezalo, 2014), 54 species of Coleoptera (22 species of Cerambycidae and 32 species of Scarabaeoidea), 101 species of Lepidoptera (Tot *et al.*, 2015) and 169 species of Heteroptera (Šeat, 2017) have been recorded so far.

The collecting of hoverflies in Serbia started in the mid-20th century and covers a period of more than 60 years. Previous investigations on the diversity of hoverflies in the wider surroundings of Vlasina Lake were few and systematic studies are lacking. They include only certain genera: *Brachyopa* Meigen, 1822 (Vujić, 1991), *Cheilosia* Meigen, 1822 (Vujić, 1996), *Criorhina* Meigen, 1822 (Vujić & Milankov, 1990), *Eristalis* Latreille, 1804 (Šimić & Vujić, 1990), *Merodon* Meigen, 1803 (Vujić *et al.*, 2012), *Neoascia* Williston, 1887 (Vujić, 1990), *Scaeva* Fabricius, 1805 (Radenković *et al.*, 1995), *Syrphus* Fabricius, 1775 (Nedeljković *et al.*, 2010) and *Volucella* Geoffroy, 1764 (Nedeljković *et al.*, 2003). However, most of these data are recorded from localities outside the boundaries of the protected area, which is why they are not included in the list given below.

Vlasina is already known as a biodiversity hotspot for many organism (IBA-Important Bird Area, IPA-Important Plant Area, PBA-Prime Butterfly Area), including hoverflies (PHA-Prime Hoverfly Area) (Vujić *et al.*, 2016). As previous hoverfly studies in the Vlasina region are scarce, the aim of this study was to summarize published and new records in order to give the first species checklist of hoverflies for this area.

Materials and Methods

Research into the hoverfly fauna of Vlasina was conducted in the period 1977-2017 within the boundaries of the protected area (Fig. 1) on six localities: 1. Vlasina Lake (42.732793N 22.335378E, 1176 m a.s.l.), 2. Delnice (42.716468N 22.313565E, 1408 m a.s.l.), 3. Vrtop (42.791688N 22.374657E, 1658 m a.s.l.), 4. Vučja River (42.75363N 22.3955E, 889 m a.s.l.), 5. Veliki Čemernik (42.72663N 22.278897E, 1531 m a.s.l.), 6. Vlasina Rid (42.725815N 22.325153E, 1308 m a.s.l.) (Fig. 2). Images of the localities are presented in Fig. 3.

Specimens were collected by the standard sweep-netting method. The collected material was prepared, pinned and labelled. Identification of adults was based on external morphological features and male terminalia using a Nikon SMZ 745T stereomicroscope. The material was identified according to the following publications: Hippa *et al.* (2001), Nedeljković (2011), Speight & Sarthou (2014), Van Veen (2004), Vujić (1990) and Vujić (1992). Insect material is deposited in the Department of Biology and Ecology, University of Novi Sad (FSUNS). A map of the investigated localities was created using QGIS 2.18.13 software (Free, 2015).




Figure 1. Boundaries of Landscape of Outstanding Features Vlasina with degrees of protection (<http://www.serbiaecotour.rs/sr/zasticena-podrucja/vlasina>).




Figure 2. Map of investigated localities.



Figure 3. Images of localities: 1. Vučja River (Photo: T. Tot), 2. Delnice (Photo: M. Vujić), 3. Vlasina Rid (Photo: M. Vujić), 4. Vlasina Lake (Photo: A. Erdeš), 5. Vrtop (Photo: T. Tot), 6. Veliki Čemernik (Photo: T. Tot).

Results

The present paper gives data on 76 species of 32 genera. The registered taxa are listed below in alphabetical order.

Anasimyia lineata (Fabricius, 1787)

New data: Vlasina: Vlasina Lake, 22.06.1991, 2♀♀, leg. A. Vujić; 11.07.2017, 4♂♂, 5♀♀, leg. T. Tot, M. Vujić.

Notes: The species prefers alluvial wetlands.

Baccha elongata (Fabricius, 1775)

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Cheilosia barbata Loew, 1857

Published in Vujić (1996).

New data: Vlasina: Vučja River, 13.07.2017, 2♂♂, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed on the Balkan Peninsula.

Ch. illustrata (Harris, 1780)

New data: Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Ch. longula (Zetterstedt, 1838)

New data: Vlasina: Vučja River, 13.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Europe, but rare on the Balkan Peninsula. In Serbia the species is protected.

Ch. proxima (Zetterstedt, 1843)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed on the Balkan Peninsula.

Ch. scutellata (Fallén, 1817)

Published in Vujić (1996).

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: Widely distributed species.

Ch. soror (Zetterstedt, 1843)

Published in Vujić (1996).

Notes: The species is widely distributed on the Balkan Peninsula.

Ch. variabilis (Panzer, 1798)

Published in Vujić (1996).

Notes: One of the most abundant species of the genus *Cheilosia* Meigen, 1822 on the Balkan Peninsula.

Ch. vernalis (Fallén, 1817)

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed on the Balkan Peninsula.

Chrysogaster solstitialis (Fallén, 1817)

New data: Vlasina: Delnice, 12.07.2017, 1♂, 4♀♀, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 2♀♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♂♂, 2♀♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Ch. bicinctum (Linnaeus, 1758)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 2♂♂, 2♀♀, leg. A. Vujić; Vlasina: Veliki Čemernik, 21.07.1993, 1♀, leg. A. Vujić, 14.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 7♂♂, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 3♂♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is predominantly distributed in the mountainous regions of Serbia.

Ch. cautum (Harris, 1776)

New data: Vlasina: Vlasina Rid, 13.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Ch. elegans Loew, 1841

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 3♂♂, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Ch. fasciatum Muller, 1764

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded on higher mountains of Serbia.

Ch. festivum (Linnaeus, 1758)

New data: Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić; 21.07.1993, 3♂♂, leg. D. Radović; 14.07.2017, 4♀♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded in lower regions of Serbia.

Ch. montanum Nedeljković & Vujić, 2015

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♀♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded only on three localities (Golija, Zlatar, Vlasina) in Serbia so far.

Ch. octomaculatum Curtis, 1837

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić.

Notes: The species is distributed mainly in the mountainous regions of Serbia.

Ch. tomentosum Giglio-Tos, 1890

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded only on several localities (Kopaonik, Šar Planina, Stara Planina, Vlasina) in Serbia so far.

Ch. vernale Loew, 1841

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded in lower regions of Serbia.

Ch. verralli Collin, 1940

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is recorded on lower mountains of Serbia.

Dasysyrphus albostrigatus (Fallén, 1817)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

D. pinastri (De Geer, 1776) sensu Doczkal, 1996

New data: Vlasina: Vlasina Lake, 16.06.1988, 1♂, leg. A. Vujić, 01-05.08.1991, 1♀, leg. A. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

D. venustus (Meigen, 1822)

New data: Vlasina; Vlasina Lake, 01.06.1988, 1♀, leg. A. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Epistrophe diaphana (Zetterstedt, 1843)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

E. grossulariae (Meigen, 1822)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 4♂♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is mainly distributed in the mountainous regions of Serbia.

Episyrphus balteatus (De Geer, 1776)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 2♂♂, leg. A. Vujić, 22.06.1991, 1♂, leg. A. Vujić, 11.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 3♀♀, leg. T. Tot, M. Vujić; Vlasina, Vlasina Rid, 13.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić; Vlasina; Vrtop, 15.07.2017, 1♂, 2♀♀, leg. T. Tot, M. Vujić.

Notes: Widely distributed species.

Eristalis arbustorum Linnaeus, 1758

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, 1♀, leg. A. Vujić; Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić, 21.07.1993, 1♂, leg. D. Radović.

Notes: This species during sampling in 2017 was numerous at every locality, so it is only recorded visually.

Eristalis jugorum Egger, 1858

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

E. pertinax (Scopoli, 1763)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 21.07.1993, 2♂♂, 1♀, leg. A. Vujić, 14.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♂♂, leg. T. Tot, M. Vujić.

Notes: The species is very common and abundant in forests in Serbia.

E. similis (Fallén, 1817)

New data: Vlasina: Veliki Čemernik, 20.07.1993, 1♀, leg. D. Radović; Vlasina: Vrtop, 15.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed on the Balkan Peninsula.

E. tenax Linnaeus, 1758

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 2♂♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. D. Radović.

Notes: This species during sampling in 2017 was numerous at every locality, so it is only recorded visually.

Eupeodes corollae (Fabricius, 1794)

New data: Vlasina: Vlasina Lake, 22.06.1991, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, 2♀♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 3♀♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 9♂♂, 5♀♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 6♂♂, 4♀♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

E. flaviceps Rondani, 1857

New data: Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded in mountainous regions of Serbia with a small number of specimens.

E. lapponicus (Zetterstedt, 1838)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

E. luniger (Meigen, 1822)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić, 14.07.2017, 3♂♂ 3♀♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Leucozona lucorum (Linnaeus, 1758) (Fig. 4).

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Melangyna cincta (Fallén, 1817)

New data: Vlasina: Vlasina Lake, 01.06.1988, 2♂♂, leg. A. Vujić.

Notes: The species is widely distributed in Serbia.



Figure 4. Female of *Leucozona lucorum* (Linnaeus, 1758) (Photo: Z. Nedeljković).

Melangyna compositarum (Verrall, 1873)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 3♂♂, leg. A. Vujić; Vlasina: Veliki Čemernik, 21.07.1991, 1♂, 3♀♀, leg. D. Radović, A. Vujić, 20.07.1993, 2♂♂, 2♀♀, leg. A. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

M. guttata (Fallén, 1817)

New data: Vlasina: Veliki Čemernik, 20.07.1993, 2♂♂, leg. A. Vujić.

Notes: The species is recorded in mountainous regions of Serbia with a small number of specimens.

M. lasiophthalma (Zetterstedt, 1843)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, 1♀, leg. A. Vujić.

Notes: The species is widely distributed in Serbia.

Melanostoma mellinum (Linnaeus, 1758)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić, 22.06.1991, 1♀, leg. A. Vujić, 11.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 2♀♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

M. scalare (Fabricius, 1794)

New data: Vlasina. Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 5♂♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♂♂, 3♀♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Meliscaeva auricollis (Meigen, 1822)

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 3♂♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is mainly distributed in the mountainous regions of Serbia.

M. cinctella (Zetterstedt, 1843)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded in mountainous regions of Serbia.

Merodon aff. *cinereus*

New data: Vlasina: Veliki Čemernik, 14.07.2017, 23♂♂, 2♀♀, leg. T. Tot, M. Vujić.

Notes: Undescribed species, potentially new for science.

M. armipes Rondani, 1843

Published in Vujić *et al.*, 2012.

Notes: The species is widely distributed on the Balkan Peninsula.

M. moenium (Wiedemann in Meigen, 1822)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 10♂♂, 1♀, leg. T. Tot, M. Vujić.

Notes: One of the most abundant species of the genus *Merodon* Meigen, 1803 on the Balkan Peninsula.

Myathropa florea (Linnaeus, 1758)

New data: Vlasina: Veliki Čemernik, 21.07.1993, 1♀, leg. D. Radović; Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Vlasina: Vlasina 13.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: Widely distributed species.

Myolepta dubia (Fabricius, 1805)

New data: Vlasina: Veliki Čemernik, 21.07.1993, 1♂, leg. A. Vujić.

Notes: The species is recorded on several mountains in Serbia with a relatively small number of specimens.

Neoascia obliqua Coe, 1940

Published in Vujić (1990).

Notes: The species is recorded in mountainous regions of Serbia.

N. tenur (Harris, 1780)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is recorded in mountainous regions and wetlands of Serbia with a small number of specimens.

Paragus haemorrhous Meigen, 1822

New data: Vlasina: Vlasina Lake, 11.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Parasyrphus lineolus (Zetterstedt, 1843)

New data: Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

P. punctulatus (Verrall, 1873)

New data: Vlasina: Vlasina Lake, 01.06.1988, 1♂, leg. A. Vujić.

Notes: The species is widely distributed in Serbia.

P. vittiger (Zetterstedt, 1843)

New data: Vlasina: Vlasina Lake, 04.07.1977, 1♀, leg. S. Šimić.

Notes: The species is distributed in the mountainous regions of Serbia.

Platycheirus albimanus (Fabricius, 1781)

New data: Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

P. europaeus Goeldlin, Maibach & Speight, 1990

New data: Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Scaeva dignota (Rondani, 1857)

Published in Radenković *et al.*, 1995.

Notes: The species is distributed in the mountainous regions of Serbia.

Scaeva pyrastris (Linnaeus, 1758)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 2♀♀, leg. A. Vujić; Veliki Čemernik, 20.07.1993, 1♂, leg. D. Radović, 14.07.2017, 7♀♀, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 4♀♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 4♀♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina: Vrtop, 15.07.2017, 9♀♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Scaeva selenitica (Meigen, 1822)

New data: Vlasina: Vlasina Lake, 22.06.1991, 1♀, leg. A. Vujić, 11.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 4♂♂, leg. T. Tot, M. Vujić; Vlasina: Vlasina: Vlasina Rid, 13.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Veliki Čemernik, 14.07.2017, 7♂♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 3♀♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Sericomyia lappona (Linnaeus, 1758)

New data: Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded only on several localities (Golija, Dubašnica, Kopaonik, Vlasina) in Serbia so far.

Spazigaster ambulans (Fabricius, 1798)

New data: Vlasina: Veliki Čemernik, 20.07.1993, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is recorded in mountainous regions of Serbia.

Sphaerophoria scripta (Linnaeus, 1758)

New data: Vlasina: Vlasina Lake, 01-05.08.1991, 1♂, leg. A. Vujić, 22.06.1991, 3♂♂, 1♀, leg. A. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♀, leg. T. Tot, M. Vujić.

Notes: Widely distributed species.

Sphegina clavata (Scopoli, 1763)

New data: Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

S. elegans Schummel, 1843

New data: Vlasina: Delnice, 12.07.2017, 2♀♀, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 4♀♀, leg. T. Tot, M. Vujić.

Notes: The species is recorded only on several localities in Serbia with a small number of specimens.

Syritta pipiens (Linnaeus, 1758)

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 2♂♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: Widely distributed species.

Syrphus ribesii (Linnaeus, 1758)

Published in Nedeljković *et al.*, 2010.

New data: Vlasina: Veliki Čemernik, 21.07.1991, 1♂♂, 2♀, leg. D. Radović, A. Vujić, 14.07.2017, 3♂♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 3♂♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina: Vrtop, 15.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

S. torvus Osten Sacken, 1875

Published in Nedeljković *et al.*, 2010.

New data: Vlasina: Vlasina Rid, 13.07.2017, 2♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 9♂♂, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♂♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

S. vitripennis Meigen, 1822

New data: Vlasina: Vlasina Lake, 01.06.1988, 1♂, 2♀, leg. A. Vujić, 01-05.08.1991, 1♂, 1♀, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 8♂♂, 3♀, leg. T. Tot, M. Vujić; Vlasina: Vlasina Rid, 13.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 5♂♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Volucella bombylans (Linnaeus, 1758)

New data: Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 1♂, 1♀, leg. T. Tot, M. Vujić.

Notes: The species is mainly distributed in the mountainous regions of Serbia.

V. pellucens (Linnaeus, 1758)

Published in Nedeljković *et al.*, 2003.

New data: Vlasina: Vlasina Lake, 11.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Delnice, 12.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 4♀♀, leg. T. Tot, M. Vujić.

Notes: The species is mainly distributed in the mountainous regions of Serbia.

V. zonaria (Poda, 1761)

New data: Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Xanthandrus comtus (Harris, 1780)

New data: Vlasina: Vlasina Lake, 01.06.1988, 1♀, leg. A. Vujić, 01-05.08.1991, 1♂, leg. A. Vujić; Vlasina: Delnice, 12.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić; Vlasina: Veliki Čemernik, 14.07.2017, 1♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 3♂♂, 4♀♀, leg. T. Tot, M. Vujić.

Notes: The species is distributed in the mountainous regions of Serbia.

Xanthogramma stackelbergi Violovitsh, 1975

New data: Vlasina: Vučja River, 13.07.2017, 1♂, leg. T. Tot, M. Vujić.

Notes: The species is mainly distributed in the mountainous regions of Serbia.

Xylota sylvarum (Linnaeus, 1758)

New data: Vlasina, Vlasina Rid, 13.07.2017, 2♀♀, leg. T. Tot, M. Vujić; Vlasina: Vrtop, 15.07.2017, 2♀♀, leg. T. Tot, M. Vujić.

Notes: The species is widely distributed in Serbia.

Discussion

Based on published literature and new data from a period of 40 years, 76 hoverfly species of 32 genera were confirmed for the Landscape of Outstanding Features "Vlasina". After true bugs (with 169 recorded species) and butterflies (with 101 recorded species), hoverflies represent one of the richest insect groups in this landscape.

The most remarkable finding is the potentially new hoverfly species for science, the description of which is in preparation: *Merodon* aff. *cinereus* (Vujić *et al.*, in prep.).

New findings of *Chrysotoxum montanum* Nedeljković & Vujić, 2015 (Nedeljković *et al.*, 2015), and *Chrysotoxum tomentosum* Giglio-Tos, 1890 (Nedeljković *et al.*, 2013), also represent very important data, since up to now *Ch. tomentosum* was collected only on Mts. Kopaonik, Šar Planina and Stara Planina, while *Ch. montanum* on Mts. Golija and Zlatar.

Although widespread in northern Europe and Siberia, *Cheilosia longula* is rare on the Balkan Peninsula. This species is only recorded on Mt. Kopaonik so far. It is listed as a protected species in Serbia based on the Code of regulations on the declaration and protection of strictly protected and protected wild species of plants, animals and fungi (Official Gazette of RS, nos. 5/2010 and 47/2011).

The richest localities, with a presence of more than 30 different species, are Vlasina Lake (38 species), Veliki Čemernik (37 species) and Vrtop (33 species). Delnice (29 species) and Vlasina Rid (19 species) are characterized by a lower number of hoverfly species, while Vučja River (16 species) is the poorest locality as regards number of species.

The most common genera were *Chrysotoxum* Meigen, 1822 (10) and *Cheilosia* Meigen, 1822 (8), while the most abundant species were *Chrysotoxum binctum*, *Chrysotoxum elegans*, *Episyrphus balteatus*, *Eristalis arbustorum*, *Eristalis tenax* and *Eupeodes corollae*, which are found at all investigated localities.

Vlasina Lake and its immediate surroundings are characterized by the presence of a specific biogeocenosis, mainly mires, wetlands, forests, meadows and pastures (according to the Law on Environmental Protection,

Official Gazette of Republic of Serbia, No.66/91). This explains why the above mentioned species are registered in this area, since more than half of them exclusively inhabit humid habitats (Speight, 2016).

Acknowledgements

The authors are thankful to the authority of the Protected Landscape Vlasina "PE Surdulica Municipality Land Development Directorate" for technical support and accommodation. Also, thanks to the members of the Scientific Research Society of Biology Students "Josif Pančić" Novi Sad. Financial support was provided by Ministry of Education, Science and Technological Development of Republic of Serbia (project OI173002).

References

- Free, Q. Q. A. (2015). Open Source Geographic Information System. Available from: <http://www.qgis.org/en/site>.
- Hippa, H., Nielsen, T. R. & Steenis, J. V. (2001). The West Palaearctic species of the genus *Eristalis* Latreille (Diptera, Syrphidae). *Norwegian Journal of Entomology*, 48, 289-327.
- Nedeljković, Z. (2011). *Taksonomska analiza vrsta iz podfamilije Syrphinae (Diptera: Syrphidae) u Srbiji*. Doktorska disertacija. Prirodno-matematički fakultet, Univerzitet u Novom Sadu, 247 pp.
- Nedeljković, Z., Vujić, A., Radenković, S. & Šimić, S. (2003). The genus *Volucella* Geoffroy, 1764 (Diptera: Syrphidae) on the Balkan Peninsula. *Acta entomologica serbica*, 8(1-2), 41-55.
- Nedeljković, Z., Vujić, A., Ricarte, A., Radenković, S. & Šimić, S. (2010). New data on the genus *Syrphus* Fabricius, 1775 (Diptera: Syrphidae) from the Balkan Peninsula including the first record of *Syrphus nitidifrons* Becker, 1921. *Acta entomologica serbica*, 15(1), 91-105.
- Nedeljković, Z., Ačanski, J., Vujić, A., Obreht, D., Đan, M., Ståhls, G. & Radenković, S. (2013). Taxonomy of *Chrysotoxum festivum* Linnaeus, 1758 (Diptera: Syrphidae)—an integrative approach. *Zoological Journal of the Linnean Society*, 169(1), 84-102.
- Nedeljković, Z., Ačanski, J., Đan, M., Obreht-Vidaković, D., Ricarte, A. & Vujić, A. (2015). An integrated approach to delimiting species borders in the genus *Chrysotoxum* Meigen, 1803 (Diptera: Syrphidae), with description of two new species. *Contribution to Zoology*, 84(4), 285-304.
- Pavićević, D. & Zatezalo, A. (2014). Insekti. In Belij, S. (Ed.), VLASINA: predeo izuzetnih odlika. *Zavod za zaštitu prirode Srbije*, Beograd, 155 pp.
- Radenković, S., Šimić, S. & Vujić, A. (1995). Genus *Scaeva* Fabricius (Diptera, Syrphidae) on the Balkan Peninsula. *Zbornik Matice srpske za prirodne nauke*, 88, 51-57.
- Rotheray, G. E. & Gilbert, F. (2011). *The natural history of hoverflies*. Forest Text, Ceredigion, UK, 334 pp.
- Speight, M. C. D. (2016). Species accounts of European Syrphidae (Diptera) 2016. *Syrph the Net, the database of European Syrphidae*, Syrph the Net publications, Dublin, 93, 288 pp.
- Speight, M. C. D. & Sarthou, J. P. (2014). StN key for the identification of the genera of European Syrphidae (Diptera). *Syrph the Net, the database of European Syrphidae*, Syrph the Net publication, Dublin, 79, 42 pp.
- Šeat, J. (2017). Prilog poznavanju stenica (Insecta: Heteroptera) Vlasine [Contribution to the knowledge on true bugs (Insecta: Heteroptera) of Vlasina]. *Zaštita prirode* 67(1-2), 43-54. [In Serbian with English abstract].
- Šimić, S. & Vujić, A. (1990). Vrste roda *Eristalis* Latreille, 1804 (Diptera Syrphidae) iz zbirke Instituta za biologiju u Novom Sadu. *Glasnik Prirodnjačkog muzeja u Beogradu*, B45, 115-126.
- Thompson, F. C. (2013). Syrphidae. *Systema Dipterorum*, Version 1.5. 13354 records. Available on: <http://www.diptera.org> [accessed on 5 September 2013].

- Tot, I. S., Slacki, A., Đurić, M. & Popović, M. (2015). Butterflies of the Vlasina region in Southeast Serbia (Lepidoptera: Papilionoidea). *Acta entomologica serbica*, 20, 117-135.
- Van Veen, M. P. (2004). *Hoverflies of Northwest Europe: identification keys to the Syrphidae*. Utrecht: KNNV Publishing, 256 pp.
- Vujić, A. (1990). Rodovi *Neoascia* Williston, 1886 i *Sphegina* Meigen, 1822 (Diptera: Syrphidae) u Jugoslaviji i opis vrste *Sphegina sublatifrons* sp. nova. *Glasnik prirodnjačkog muzeja u Beogradu*, B45, 77-93.
- Vujić, A. (1991). Vrste roda *Brachyopa* Meigen 1822 (Diptera: Syrphidae) u Jugoslaviji. *Glasnik prirodnjačkog muzeja u Beogradu*, B46, 141-150.
- Vujić, A. (1992). *Taksonomski položaj i zoogeografska analiza roda Cheilosia i srodnih rodova (Diptera: Syrphidae) na Balkanskom poluostrvu*. Doktorska disertacija. Prirodno-matematički fakultet. Univerzitet u Novom Sadu, 278 pp.
- Vujić, A. (1996). *Genus Cheilosia Meigen and related genera (Diptera: Syrphidae) on the Balkan Peninsula: Rod Cheilosia Meigen i srodni rodovi (Diptera: Syrphidae) na Balkanskom poluostrvu*. Matica srpska, Novi Sad, 194 pp.
- Vujić, A. & Milankov, V. (1990). Taksonomski status vrsta roda *Criorrhina* Meigen 1822 (Diptera: Syrphidae) zabeleženih u Jugoslaviji. *Glasnik prirodnjačkog muzeja u Beogradu*, B45, 105-114.
- Vujić, A., Radenković, S., Stähls, G., Ačanski, J., Stefanović, A., Veselić, S., Andrić, A. & Hayat, R. (2012). Systematics and taxonomy of the *ruficornis* group of genus *Merodon* Meigen (Diptera: Syrphidae). *Systematic Entomology*, 37(3), 578-602.
- Vujić, A., Radenković, S., Nikolić, T., Radišić, D., Trifunov, S., Andrić, A., Markov, Z., Jovičić, S., Mudri Stojnić, S., Janković, M. & Lugonja, P. (2016). Prime Hoverfly (Insecta: Diptera: Syrphidae) Areas (PHA) as a conservation tool in Serbia. *Biological Conservation*, 198, 22-32.

ФАУНА ОСОЛИКИХ МУВА (DIPTERA: SYRPHIDAE) ПРЕДЕЛА ИЗУЗЕТНИХ ОДЛИКА “ВЛАСИНА”

ТАМАРА ТОТ, МИХАИЛО ВУЈИЋ, ЛАУРА ЛИКОВ, ЗОРИЦА НЕДЕЉКОВИЋ,
СНЕЖАНА РАДЕНКОВИЋ И АНТЕ ВУЈИЋ

ИЗВОД

Током вишегодишњих теренских истраживања организованих у периоду од 1977-2017. године на шест локалитета унутар подручја Предео изузетних одлика “Власина” (југоисточна Србија) сакупљено је укупно 76 врста осоликих мува. Међу идентификованим таксонима налази се ретка врста *Cheilosis longula* (Zetterstedt, 1838), која је заштићена у Србији. Поред тога забележена је и једна потенцијално нова врста за науку (*Merodon* aff. *cinereus*). У раду је дата прва листа осоликих мува за ово подручје.

Received: January 1st, 2018.

Accepted: May 21st, 2018.